

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

I declare this is my own work.

GCSE CHEMISTRY

H

Higher Tier Paper 1

Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- a scientific calculator
- the periodic table (enclosed).

Instructions

- Use black ink or black ball-point pen.
- Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

Information

- The maximum mark for this paper is 100.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	



0 1

This question is about carbon and its compounds.

Fullerenes are molecules of carbon atoms.

The first fullerene to be discovered was Buckminsterfullerene (C_{60}).

0 1 . 1

What shape is a Buckminsterfullerene molecule?

[1 mark]

0 1 . 2


Give **one** use of a fullerene.

[1 mark]

Propanone is a compound of carbon, hydrogen and oxygen.

Figure 1 shows the dot and cross diagram for a propanone molecule.

Figure 1




0 1 . 3 Complete **Figure 2** to show a propanone molecule.

Use a line to represent each single bond.

Use **Figure 1**.

[1 mark]

Figure 2



0 1 . 4 Determine the molecular formula of propanone.

Use **Figure 1**.

[1 mark]

Molecular formula = _____

0 1 . 5 Propanone is a liquid with a low boiling point.

Why does propanone have a low boiling point?

[1 mark]

Tick (✓) **one** box.

The covalent bonds are strong.

☐

The covalent bonds are weak.

☐

The intermolecular forces are strong.

☐

The intermolecular forces are weak.

☐

Turn over ►



Turn over for the next question

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ►



0 2

This question is about atomic structure and the periodic table.

Gallium (Ga) is an element that has two isotopes.

0 2 . 1

Give the meaning of 'isotopes'.

You should answer in terms of subatomic particles.

[2 marks]

0 2 . 2

Table 1 shows the mass numbers and percentage abundances of the isotopes of gallium.

Table 1

Mass number	Percentage abundance (%)
69	60
71	40

Calculate the relative atomic mass (A_r) of gallium.

Give your answer to 1 decimal place.

[2 marks]

Relative atomic mass (1 decimal place) = _____



Gallium (Ga) is in Group 3 of the modern periodic table.

0 2 . 3

Give the numbers of electrons and neutrons in an atom of the isotope $^{69}_{31}\text{Ga}$

[2 marks]

Number of electrons _____

Number of neutrons _____

0 2 . 4

What is the most likely formula of a gallium ion?

[1 mark]

Tick (✓) **one** box.

Ga⁺ ☐

Ga⁻ ☐

Ga³⁺ ☐

Ga³⁻ ☐

0 2 . 5

Gallium was discovered six years after Mendeleev published his periodic table.

Give **two** reasons why the discovery of gallium helped Mendeleev's periodic table to become accepted.

[2 marks]

1 _____

2 _____

9

Turn over ►




0 3

This question is about the extraction of metals.

Element **R** is extracted from its oxide by reduction with hydrogen.

The equation for the reaction is:

**0 3****. 1**

The sum of the relative formula masses (M_r) of the reactants ($3\text{H}_2 + \text{RO}_3$) is 150

Calculate the relative atomic mass (A_r) of **R**.

Relative atomic masses (A_r): H = 1 O = 16

[2 marks]

Relative atomic mass (A_r) of **R** = _____

0 3**. 2**

Identify element **R**.

You should use:

- your answer to **question 03.1**
- the periodic table.

[1 mark]


Identity of **R** = _____



0	3	.	3
---	---	---	---

Carbon is used to extract tin (Sn) from tin oxide (SnO₂).

The equation for the reaction is:



Calculate the percentage atom economy for extracting tin in this reaction.

Relative atomic masses (*A_r*): C = 12 O = 16 Sn = 119

[3 marks]

Percentage atom economy = _____ %

Question 3 continues on the next page

Turn over ►



Tungsten is extracted from tungsten oxide (WO_3).

Table 2 shows information about three possible methods to extract tungsten from tungsten oxide.

Table 2

Method	Reactant	Relative cost of reactant	Products
1	Carbon	Low	Tungsten solid Carbon dioxide gas Tungsten carbide solid
2	Hydrogen	High	Tungsten solid Water vapour
3	Iron	Low	Tungsten solid Iron oxide solid

Evaluate the three possible methods for extracting tungsten from tungsten oxide.

[4 marks]

[illegible]

10



0	4
---	---

This question is about Group 1 elements.

0	4	.	1
---	---	---	---

Give **two** observations you could make when a small piece of potassium is added to water.

[2 marks]

1 _____

2 _____

0	4	.	2
---	---	---	---

Complete the equation for the reaction of potassium with water.

You should balance the equation.

[2 marks]



0	4	.	3
---	---	---	---

Explain why the reactivity of elements changes going down Group 1.

[4 marks]

Turn over ►



Sodium reacts with oxygen to produce the ionic compound sodium oxide.

Oxygen is a Group 6 element.

0	4	.	4
---	---	---	---

Draw a dot and cross diagram to show what happens when atoms of sodium and oxygen react to produce sodium oxide.

[4 marks]

Diagram



0 4 . 5

Why is oxygen described as being reduced in the reaction between sodium and oxygen?

[1 mark]

0 4 . 6

Explain why sodium oxide has a high melting point.

[3 marks]

16

Turn over for the next question

Turn over ►



0 5

This question is about salts.

0 5 . 1

Name the salt produced by the neutralisation of hydrochloric acid with potassium hydroxide.

[1 mark]

0 5 . 2

Write an ionic equation for the neutralisation of hydrochloric acid with potassium hydroxide.

[1 mark]

_____ + _____ → _____

0 5 . 3

Soluble salts can be produced by reacting dilute hydrochloric acid with an insoluble solid.

Copper, copper carbonate and copper oxide are insoluble solids.

Which of these insoluble solids can be used to make a copper salt by reacting the solid with dilute hydrochloric acid?

[1 mark]Tick (✓) **one** box.

Copper and copper carbonate only

☐

Copper and copper oxide only

☐

Copper carbonate and copper oxide only

☐

Copper, copper carbonate and copper oxide

☐

A student makes crystals of magnesium sulfate.

This is the method used.

1. Add sulfuric acid to a beaker.
2. Warm the sulfuric acid.
3. Add a spatula of magnesium oxide to the beaker.
4. Stir the mixture.
5. Repeat steps 3 and 4 until there is magnesium oxide remaining in the beaker.
6. Filter the mixture.
7. Evaporate the filtrate gently until crystals start to form.
8. Leave the solution to finish crystallising.

0 5 . 4

Give **one** reason for:

- step 2
- step 5
- step 6.

[3 marks]

Step 2 _____

Step 5 _____

Step 6 _____

0 5 . 5

How should the filtrate be evaporated gently in **step 7**?

[1 mark]


Turn over ►



0 5 . 6

Iron chloride is produced by heating iron in chlorine gas.

The equation for the reaction is:



Calculate the volume of chlorine needed to react with 14 g of iron.

You should calculate:

- the number of moles of iron used
- the number of moles of chlorine that react with 14 g of iron
- the volume of chlorine needed.

Relative atomic mass (A_r): Fe = 56

The volume of 1 mole of gas = 24 dm³

[3 marks]

Volume of chlorine = _____ dm³

10



0 6

This question is about metals.

0 6 . 1**Table 3** shows information about four substances.**Table 3**

Substance	Melting point in °C	Boiling point in °C	Does it conduct electricity in the solid state?	Does it conduct electricity in the liquid state?
A	–117	79	No	No
B	801	1413	No	Yes
C	1535	2750	Yes	Yes
D	1610	2230	No	No

Which substance could be a metal?

[1 mark]Tick (✓) **one** box.**A**☐**B**☐**C**☐**D**☐**0 6 . 2**

Explain why alloys are harder than pure metals.

[3 marks]

Turn over ►

0 6 . 3

A student wants to compare the reactivity of an unknown metal, **Q**, with that of zinc.

Both metals are more reactive than silver.

The student is provided with:

- silver nitrate solution
- metal **Q** powder
- zinc powder
- a thermometer
- normal laboratory equipment.

No other chemicals are available.

Describe a method the student could use to compare the reactivity of metal **Q** with that of zinc.

Your method should give valid results.

[4 marks]

8



07

This question is about chemical reactions and electricity.

07.1

Electrolysis and chemical cells both involve chemical reactions and electricity.

Explain the difference between the processes in electrolysis and in a chemical cell.

[2 marks]

07.2


A teacher demonstrates the electrolysis of molten lead bromide.

Bromine is produced at the positive electrode.

Complete the half equation for the production of bromine.

You should balance the half equation.

[2 marks]



07.3

Two aqueous salt solutions are electrolysed using inert electrodes.

Complete **Table 4** to show the product at each electrode.

[3 marks]

Table 4


Salt solution	Product at positive electrode	Product at negative electrode
Copper nitrate		copper
Potassium iodide		

Turn over ►



Some students investigated the electrolysis of copper nitrate solution using inert electrodes.

Figure 4 shows the apparatus.



The students investigated how the mass of copper produced at the negative electrode varied with:

- time
- current.

This is the method used.

1. Weigh the negative electrode.
2. Set up the apparatus shown in **Figure 4**.
3. Adjust the power supply until the ammeter shows a current of 0.3 A
4. Switch off the power supply after 5 minutes.
5. Rinse the negative electrode with water and allow to dry.
6. Reweigh the negative electrode.
7. Repeat steps 1 to 6 for different times.
8. Repeat steps 1 to 7 at different currents.

0	7	.	4
---	---	---	---

Some of the copper produced did not stick to the negative electrode but fell to the bottom of the beaker.

Suggest how the students could find the total mass of copper produced.

[4 marks]


Question 7 continues on the next page

Turn over ►



The students plotted their results on a graph.

Figure 5 shows the graph.



A student correctly concluded that the total mass of copper produced is directly proportional both to the time and to the current.

0 7 . 5

How do the results in **Figure 5** support the conclusion that the total mass of copper produced is directly proportional to the time?

[1 mark]

0 7 . 6

How do the results in **Figure 5** support the conclusion that the total mass of copper produced is directly proportional to the current?

Use data from **Figure 5** in your answer.

[1 mark]



0 7 . 7

Copper nitrate solution is blue.

Suggest why the blue colour of the copper nitrate solution fades during the electrolysis.

[1 mark]

0 7 . 8

Determine the number of atoms of copper produced when copper nitrate solution is electrolysed for 20 minutes at a current of 0.6 A

Give your answer to 3 significant figures.

Use **Figure 5**.

Relative atomic mass (A_r): Cu = 63.5

The Avogadro constant = 6.02×10^{23} per mole

[3 marks]

Number of atoms (3 significant figures) = _____

17

Turn over for the next question


Turn over ►



0 8

This question is about the reaction between hydrogen sulfide (H_2S) and oxygen.

The equation for the reaction is:



0 8 . 1

What does $\text{H}_2\text{O}(\text{g})$ represent?

[1 mark]

0 8 . 2

Calculate the volume of oxygen required to react with 50 cm^3 of hydrogen sulfide.

[1 mark]

Volume = _____ cm^3

0 8 . 3

Figure 6 shows part of the reaction profile for the reaction.

The reaction is exothermic.

Complete **Figure 6**.

You should:

- complete the profile line
- label the activation energy
- label the overall energy change.

[3 marks]

Figure 6




Figure 7 shows the displayed formula equation for the reaction of hydrogen sulfide with oxygen.

$$2\text{H}-\text{S}-\text{H} + 3\text{O}=\text{O} \longrightarrow 2\text{H}-\text{O}-\text{H} + 2\text{O}=\text{S}=\text{O}$$

Table 5

Bond	H—S	O=O	H—O	S=O
Energy in kJ/mol	364	498	464	X

Calculate the bond energy **X** for the $\text{S}=\text{O}$ bond.

[5 marks]

[illegible]

X = kJ/mol

10

Turn over ►



0 9

This question is about acids.

Hydrogen chloride and ethanoic acid both dissolve in water.

All hydrogen chloride molecules ionise in water.

Approximately 1% of ethanoic acid molecules ionise in water.

0 9 . 1

A solution is made by dissolving 1 g of hydrogen chloride in 1 dm³ of water.

Which is the correct description of this solution?

[1 mark]

Tick (✓) **one** box.

A concentrated solution of a strong acid

☐

A concentrated solution of a weak acid

☐

A dilute solution of a strong acid

☐

A dilute solution of a weak acid

☐**0 9 . 2**

Which solution would have the lowest pH?

[1 mark]

Tick (✓) **one** box.

0.1 mol/dm³ ethanoic acid solution

☐

0.1 mol/dm³ hydrogen chloride solution

☐

1.0 mol/dm³ ethanoic acid solution

☐

1.0 mol/dm³ hydrogen chloride solution

☐

A student investigated the concentration of a solution of sodium hydroxide by titration with a 0.0480 mol/dm^3 ethanedioic acid solution.

This is the method used.

1. Measure 25.0 cm^3 of the sodium hydroxide solution into a conical flask using a 25.0 cm^3 pipette.
2. Add two drops of indicator to the sodium hydroxide solution.
3. Fill a burette with the 0.0480 mol/dm^3 ethanedioic acid solution to the 0.00 cm^3 mark.
4. Add the ethanedioic acid solution to the sodium hydroxide solution until the indicator changes colour.
5. Read the burette to find the volume of the ethanedioic acid solution used.

0 9 . 3

Suggest **two** improvements to the method that would increase the accuracy of the result.

[2 marks]

1 _____

2 _____

Question 9 continues on the next page

Turn over ►



0 9 . 4

Ethanedioic acid is a solid at room temperature.

Calculate the mass of ethanedioic acid ($\text{H}_2\text{C}_2\text{O}_4$) needed to make 250 cm^3 of a solution with concentration 0.0480 mol/dm^3

Relative formula mass (M_r): $\text{H}_2\text{C}_2\text{O}_4 = 90$


[2 marks]

Mass = _____ g

0 9 . 5

The student found that 25.0 cm^3 of the sodium hydroxide solution was neutralised by 15.00 cm^3 of the 0.0480 mol/dm^3 ethanedioic acid solution.

The equation for the reaction is:



Calculate the concentration of the sodium hydroxide solution in mol/dm^3

[3 marks]

Concentration = _____ mol/dm^3

9

END OF QUESTIONS



There are no questions printed on this page


*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

[illegible]



**GCSE
CHEMISTRY
8462/1H**

Paper 1 Higher Tier

Mark scheme

June 2021

Version: 1.0 Final Mark Scheme



Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from aqa.org.uk

Copyright information

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2021 AQA and its licensors. All rights reserved.

Information to Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the Examiner make his or her judgement
- the Assessment Objectives and specification content that each question is intended to cover.

The extra information is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

2. Emboldening and underlining

- 2.1** In a list of acceptable answers where more than one mark is available 'any **two** from' is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **or**. Different terms in the mark scheme are shown by a / ; eg allow smooth / free movement.
- 2.4** Any wording that is underlined is essential for the marking point to be awarded.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general principle to be followed in such a situation is that 'right + wrong = wrong'.

Each error / contradiction negates each correct response. So, if the number of error / contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (indicated as * in example 1) are not penalised.

Example 1: What is the pH of an acidic solution?

[1 mark]

Student	Response	Marks awarded
1	green, 5	0
2	red*, 5	1
3	red*, 8	0

Example 2: Name two planets in the solar system.

[2 marks]

Student	Response	Marks awarded
1	Neptune, Mars, Moon	1
2	Neptune, Sun, Mars, Moon	0

3.2 Use of chemical symbols / formulae

If a student writes a chemical symbol / formula instead of a required chemical name, full credit can be given if the symbol / formula is correct and if, in the context of the question, such action is appropriate.

3.3 Marking procedure for calculations

Marks should be awarded for each stage of the calculation completed correctly, as students are instructed to show their working. Full marks can, however, be given for a correct numerical answer, without any working shown.

3.4 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.5 Errors carried forward

Any error in the answers to a structured question should be penalised once only.

Papers should be constructed in such a way that the number of times errors can be carried forward is kept to a minimum. Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ecf in the marking scheme.

3.6 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term.

3.7 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.8 Allow

In the mark scheme additional information, 'allow' is used to indicate creditworthy alternative answers.

3.9 Ignore

Ignore is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

3.10 Do not accept

Do **not** accept means that this is a wrong answer which, even if the correct answer is given as well, will still mean that the mark is not awarded.

4. Level of response marking instructions

Extended response questions are marked on level of response mark schemes.

- Level of response mark schemes are broken down into levels, each of which has a descriptor.
- The descriptor for the level shows the average performance for the level.
- There are two marks in each level.

Before you apply the mark scheme to a student's answer, read through the answer and annotate it (as instructed) to show the qualities that are being looked for. You can then apply the mark scheme.

Step 1: Determine a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the descriptor for that level. The descriptor for the level indicates the different qualities that might be seen in the student's answer for that level. If it meets the lowest level then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer.

When assigning a level you should look at the overall quality of the answer. Do **not** look to penalise small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level.

Use the variability of the response to help decide the mark within the level, ie if the response is predominantly level 2 with a small amount of level 3 material it would be placed in level 2 but be awarded a mark near the top of the level because of the level 3 content.

Step 2: Determine a mark

Once you have assigned a level you need to decide on the mark. The descriptors on how to allocate marks can help with this.

The exemplar materials used during standardisation will help. There will be an answer in the standardising materials which will correspond with each level of the mark scheme. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the example to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the example.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.

Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other valid points. Students do **not** have to cover all of the points mentioned in the indicative content to reach the highest level of the mark scheme.

You should ignore any irrelevant points made. However, full marks can be awarded only if there are no incorrect statements that contradict a correct response.

An answer which contains nothing of relevance to the question must be awarded no marks.

Question 1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.1	spherical	allow ball-shaped ignore round / circular	1	AO1 4.2.3.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.2	any one from: <ul style="list-style-type: none"> • drug delivery (round the body) • hydrogen storage • anti-oxidants • reduction of bacterial growth • catalysts • (cylindrical fullerenes for) strengthening materials • (spherical fullerenes for) lubricants 		1	AO1 4.2.3.3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.3	$ \begin{array}{ccccccc} & & \text{H} & & & \text{H} & \\ & & & & & & \\ \text{H} & - & \text{C} & - & \text{C} & - & \text{C} & - & \text{H} \\ & & & & & & & \\ & & \text{H} & & \text{O} & & \text{H} & \end{array} $		1	AO2 4.2.1.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.4	C ₃ H ₆ O	allow CH ₃ COCH ₃ allow elements in any order	1	AO2 4.2.1.4

Question 1 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
01.5	the intermolecular forces are weak		1	AO1 4.2.2.1 4.2.2.4

Question	Answers	Mark	AO/ Spec. Ref
01.6	Level 3: Relevant points (reasons/causes) are identified, given in detail and logically linked to form a clear account.	5–6	AO1 4.2.2.6 4.2.3.2
	Level 2: Relevant points (reasons/causes) are identified, and there are attempts at logical linking. The resulting account is not fully clear.	3–4	
	Level 1: Points are identified and stated simply, but their relevance is not clear and there is no attempt at logical linking.	1–2	
	No relevant content	0	
	Indicative content <ul style="list-style-type: none"> • bonds are covalent • giant / macromolecular structure • three (covalent) bonds per carbon atom or only three electrons per carbon atom used in (covalent) bonds • so one electron per carbon atom (is delocalised) • these delocalised electrons • can move through the structure • carrying (electrical) charge • so graphite conducts electricity • layered structure • of (interlocking) hexagonal rings • with weak (intermolecular) forces between layers or no (covalent) bonds between layers • so the layers can slide over each other • so graphite is soft and slippery 		

Total			11
--------------	--	--	-----------

Question 2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.1	(atoms with the) same number of protons	allow atoms with the same atomic number allow atoms of the same element	1	AO1 4.1.1.4 4.1.1.5
	(but with) different numbers of neutrons	ignore the same number of electrons ignore (but with) different mass numbers do not accept (but with) different relative atomic mass	1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.2	$(A_r =) \frac{(69 \times 60) + (71 \times 40)}{100}$		1	AO2 4.1.1.6
	= 69.8		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.3	(number of electrons) = 31		1	AO2 4.1.1.4 4.1.1.5
	(number of neutrons) = 38		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.4	Ga^{3+}		1	AO3 4.2.1.2

Question 2 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
02.5	(gallium) fitted in a gap (Mendeleev had left)		1	AO2 4.1.2.2
	(gallium's) properties were predicted correctly (by Mendeleev)	allow (gallium's) properties matched the rest of the group	1	
Total			9	

Question 3

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.1	$(3 \times M_r \text{ H}_2\text{O} = 3 \times (2 + 16) =) 54$		1	AO2
	$(A_r \text{ R} = 150 - 54 =) 96$	ignore units	1	4.3.1.1
	alternative approach: $(M_r \text{ RO}_3 = 150 - 6 =) 144 (1)$			4.3.1.2
	$(A_r \text{ R} = 144 - (3 \times 16) =) 96 (1)$	ignore units		

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.2	(R =) molybdenum / Mo	allow ecf from question 03.1	1	AO3 4.1.1.1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
03.3	(total M_r of reactants) = 163		1	AO2
	(% atom economy =) $\frac{119}{163} (\times 100)$	allow correct use of an incorrectly calculated value of total M_r	1	4.3.1.2
	= 73 (%)	allow 73.00613 (%) correctly rounded to at least 2 significant figures	1	4.3.3.2

Question 3 continued

Question	Answers	Mark	AO/ Spec. Ref
03.4	Level 2: Some logically linked reasons are given. There may also be a simple judgement.	3-4	AO3 4.4.1.3
	Level 1: Relevant points are made. They are not logically linked.	1-2	
	No relevant content	0	
	Indicative content <ul style="list-style-type: none"> • carbon and iron are the cheapest reactants • hydrogen is the most expensive reactant • separating solid products is expensive • separating solid products is time consuming • in method 1, tungsten needs to be separated from tungsten carbide • in method 1, some tungsten is lost as tungsten carbide • in method 1, the carbon dioxide produced will escape • in method 2, the water vapour produced will escape • in method 2, no separation of solids is needed • in method 3, tungsten needs to be separated from iron oxide 		
Total			10








Question 4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.1	any two from: <ul style="list-style-type: none"> • (potassium) floats • (potassium) melts • (potassium) moves around • potassium becomes smaller • (lilac) flame • effervescence 	allow potassium disappears allow fizzing	2	AO1 4.1.2.5 4.4.1.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.2	$2\text{K} + 2\text{H}_2\text{O} \rightarrow 2\text{KOH} + \text{H}_2$	allow multiples allow 1 mark for KOH and H ₂	2	AO1 AO2 4.1.1.1 4.1.2.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.3	reactivity increases (going down the group)		1	AO1 4.1.2.5 4.4.1.2
	(because) the outer electron / shell is further from the nucleus	allow (because) there are more shells allow (because) the atoms get larger	1	
	(so) there is less attraction between the nucleus and the outer electron / shell	allow (so) there is more shielding from the nucleus do not accept incorrect attractions	1	
	(so) the atom loses an electron more easily		1	

Question 4 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.4	(dot and cross diagram to show) sodium atom and oxygen atom	allow use of outer shells only	1	AO2 4.2.1.1 4.2.1.2
	two sodium atoms to one oxygen atom	allow two sodium ions to one oxide ion	1	
	(to produce) sodium ion with a + charge		1	
	(to produce) oxide ion with a 2- charge		1	
<div><div></div><div></div><div></div><div></div><div><div></div><div></div><div></div></div></div>				
scores 4 marks				

Question 4 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.5	(oxygen) gains electrons		1	AO1 4.4.1.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
04.6	giant structure	allow (giant ionic) lattice	1	AO1 4.2.1.3 4.2.2.1 4.2.2.3
	(with) strong (electrostatic) forces of attraction between (oppositely charged) ions		1	
	(so) large amounts of energy are needed to break the bonds / forces	allow (so) large amounts of energy are needed to separate the ions	1	

Total			16	
--------------	--	--	-----------	--

Question 5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.1	potassium chloride	allow KCl	1	AO1 4.4.2.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.2	$\text{H}^+ + \text{OH}^- \rightarrow \text{H}_2\text{O}$	ignore state symbols	1	AO1 4.1.1.1 4.4.2.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.3	copper carbonate and copper oxide only		1	AO1 4.4.1.2 4.4.2.2 4.4.2.3 RPA1

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.4	(Step 2) to speed up the reaction	ignore to remove impurities	1	AO1 4.4.2.3 RPA1
	(Step 5) to make sure all the (hydrochloric) acid reacts		1	
	(Step 6) to remove the excess magnesium oxide		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.5	using a (boiling) water bath or using an electric heater		1	AO1 4.4.2.3 RPA1

Question 5 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
05.6	(moles Fe = $\frac{14}{56}$ =) 0.25 (mol)		1	AO2 4.3.2.1 4.3.2.2 4.3.5
	(moles Cl ₂ = $\frac{3}{2}$ × 0.25 =) 0.375 (mol)	allow correct use of an incorrectly calculated number of moles of Fe	1	
	(volume Cl ₂ = 24 × 0.375) = 9.0 (dm ³)	allow correct use of an incorrectly calculated number of moles of Cl ₂	1	
Total			10	

Question 6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.1	C		1	AO3 4.2.2.7 4.2.2.8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.2	(in an alloy) the atoms are of different sizes		1	AO1 4.2.2.7
	(so) the layers (of atoms in an alloy) are distorted		1	
	(so in an alloy) the layers slide over each other less easily (than in a pure metal)		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
06.3	measure temperature change	allow measure the temperature before and after the reaction	1	AO1
	when each metal is added to silver nitrate solution		1	AO1
	same concentration / volume of solution or same mass / moles of metal	allow same initial temperature (of silver nitrate solution)	1	AO2
	the greater the temperature change the more reactive		1	AO3 4.4.1.2 4.5.1.1 RPA4

Total			8	
--------------	--	--	----------	--

Question 7

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.1		allow voltage for electricity allow potential difference for electricity allow (electrical) current for electricity		AO1 4.4.3.1 4.5.2.1
	electrolysis uses electricity to produce a chemical reaction	allow electrolysis uses electricity to decompose a compound / electrolyte	1	
	(but) cells use a chemical reaction to produce electricity		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.2	$2\text{Br}^- \rightarrow \text{Br}_2 + 2\text{e}^-$	allow multiples allow 1 mark for Br_2 and e^-	2	AO2 4.1.1.1 4.4.3.1 4.4.3.2 4.4.3.5

Question	Answers		Extra information	Mark	AO / Spec. Ref.
07.3	Salt solution	Product at positive electrode	Product at negative electrode		AO2 4.4.1.2 4.4.3.4 RPA3
	(copper nitrate)	oxygen (1)	(copper)	1	
	(potassium iodide)	iodine (1)	hydrogen (1)	2	

Question 7 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.4	filter the mixture		1	AO3 4.1.1.2 4.4.3.4 RPA3
	wash and dry the copper / residue		1	
	weigh the copper collected		1	
	add to the increase in mass of the electrode		1	

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.5	(for given current) straight line through the origin	allow (for given current) when time doubles, mass doubles	1	AO3 4.4.3.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.6	(for given time) when current doubles, mass doubles with supporting data		1	AO3 4.4.3.4

Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.7	copper ions are discharged (from the solution)	allow the solution becomes less concentrated allow copper ions are removed (from the solution) allow copper ions are used up (from the solution)	1	AO3 4.4.3.1

Question 7 continued


Question	Answers	Extra information	Mark	AO / Spec. Ref.
07.8	(number of moles = $\frac{0.24}{63.5}$ =) 3.78×10^{-3} or 0.00378		1	AO2 4.3.2.1
	(number of atoms =) $0.00378 \times 6.02 \times 10^{23}$	allow correct use of an incorrectly calculated number of moles	1	
	= 2.28×10^{21}	allow a correct evaluation to 3 significant figures of an incorrect expression which involves only a mass from the graph, the A_r of copper and the Avogadro constant	1	

Total			17
--------------	--	--	-----------

Question 8

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.1	water vapour	allow steam allow gaseous water	1	AO1 4.1.1.1 4.2.2.2

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.2	75 (cm ³)		1	AO2 4.3.5

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.3	product level below reactants	ignore labelling of products	1	AO1 4.5.1.2
	activation energy drawn and labelled		1	
	overall energy change drawn and labelled	if endothermic profile drawn allow corresponding overall energy change	1	
 <p>scores 3 marks</p>				

Question 8 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
08.4	(bonds broken = $4(364) + 3(498)$ =) 2950		1	AO2 4.5.1.3
	(bonds formed = $2950 + 1034$ =) 3984	allow correct use of incorrectly calculated values of bonds broken	1	
	$4X + 4(464) = 3984$	allow correct use of incorrectly calculated values of bonds formed	1	
	$4X = (3984 - 1856 =) 2128$		1	
	$X = 532$ (kJ/mol)		1	
	alternative approach: (bonds broken = $4(364) + 3(498)$ =) 2950 (1)			
	(bonds formed = $4(464) + 4X$ =) $1856 + 4X$ (1)			
	$(1856 + 4X) - 2950 = 1034$ (1)	allow correct use of incorrectly calculated values of bonds broken and/or bonds formed		
	$4X = (1034 + 2950 - 1856 =) 2128$ (1)			
	$X = 532$ (kJ/mol) (1)			
Total			10	

Question 9

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.1	a dilute solution of a strong acid		1	AO2 4.3.2.5 4.4.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.2	1.0 mol/dm ³ hydrogen chloride solution		1	AO2 4.4.2.4 4.4.2.6

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.3	any two from: <ul style="list-style-type: none"> swirl (the solution) white tile (under the flask) add (ethanedioic) acid dropwise (near the endpoint) repeat and calculate mean 		2	AO3 4.4.2.5 RPA2

Question 9 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.4	(concentration = $90 \times 0.0480 =$) 4.32 (g/dm ³)		1	AO2 4.3.2.1 4.3.2.5 4.3.4
	(mass = $4.32 \times \frac{250}{1000}$) = 1.08 (g) alternative approach: (moles = $0.0480 \times \frac{250}{1000} =$) 0.012 (mol) (1) (mass = 0.012×90) = 1.08 (g) (1)	allow correct use of an incorrectly calculated value of concentration in g/dm ³ allow correct use of an incorrectly calculated value of number of moles	1	

Question 9 continued

Question	Answers	Extra information	Mark	AO / Spec. Ref.
09.5	(moles $\text{H}_2\text{C}_2\text{O}_4 = \frac{15.0}{1000} \times 0.0480$) = 0.00072 (mol)		1	AO2 4.3.4 4.4.2.5 RPA2
	(moles NaOH = moles $\text{H}_2\text{C}_2\text{O}_4 \times 2 =$) 0.00144 (mol)	allow correct use of an incorrectly calculated value of number of moles of $\text{H}_2\text{C}_2\text{O}_4$	1	
	(concentration = $\frac{0.00144}{25.0} \times 1000$) = 0.0576 (mol/dm ³)	allow 0.058 (mol/dm ³) allow correct use of an incorrectly calculated value of number of moles of NaOH	1	
	alternative approach: $\frac{\text{volume} \times \text{conc (acid)}}{\text{volume} \times \text{conc (NaOH)}} = \frac{1}{2} (1)$ (conc NaOH =) $2 \times \frac{15.0 \times 0.0480}{25.0} (1)$ = 0.0576 (mol/dm ³) (1)	allow inverse allow correct use of incorrect mole ratio		

Total			9
--------------	--	--	----------